

Supersymmetric Wilson Loops in $\mathcal{N} = 6$ Super Chern-Simons-matter theory

Bin Chen

Department of Physics,
and State Key Laboratory of Nuclear Physics and Technology,
Peking University,
Beijing 100871, P.R. China
E-mail: bchen01@pku.edu.cn

Jun-Bao Wu

International School for Advanced Studies (SISSA),
via Beirut 2-4, I-34014 Trieste, Italy
and INFN, Trieste section, Via Valerio 2, I - 34127 Trieste, Italy
E-mail: wujunbao@sissa.it

September 29, 2009

Abstract

We study supersymmetric Wilson loop operators in ABJM theory from both sides of the AdS_4/CFT_3 correspondence. We first construct some supersymmetric Wilson loops. The perturbative computations are performed in the field theory side at the first two orders. A fundamental string solution ending on a circular loop is also studied.

1 Introduction

Four-dimensional $\mathcal{N} = 4$ super Yang-Mills theory is one of the most remarkable field theories. Among other things, this theory appears as low energy effective theory of multi D3-branes. This leads to the famous AdS_5/CFT_4 correspondence [1, 2, 3, 4]. Besides the local operators, supersymmetric Wilson loops play an important role in this correspondence. The vacuum expectation value (VEV) of half-BPS straight Wilson lines equals to unit due to the global supersymmetries they preserve. But the vacuum expectation value of half-BPS circular Wilson loop depends on the coupling constant in a nontrivial way. The computation of this VEV can be reduced to a computation via a Gaussian matrix model. Evidence of this reduction was found first by direct perturbative calculations [5]: in the Feynman gauge, the contributions from the diagrams which are not ladder/rainbow diagrams cancel each other. Later, this reduction was obtained from the conformal transformation linking the circular loop and the straight line [6]. Recently this reduction was obtained through localization technique [7]. For the Wilson loop in the fundamental representation of the gauge group, the VEV computed via this matrix model has the same large N , large λ behavior as the one from the string theory dual description in terms of fundamental strings in AdS_5 ending on this loop [8, 9, 10, 11]. Similar conclusion was found for Wilson loops in symmetric (anti-symmetric) representation, in which case a better dual description is in terms of $D3$ ($D5$) branes [12, 13, 14, 15, 16]. Various aspects of Wilson loops with less supersymmetries were discussed in [17]-[24].

The low energy effective theory of multi M2-branes and M5-branes is one of the most important subjects in M-theory. For multi M2-branes in flat eleven dimensional spacetime, the low energy effective theory is the IR limit of $\mathcal{N} = 8$ three dimensional super Yang-Mills theory. The construction of $\mathcal{N} = 8$ three dimensional conformal field theory for multi-membranes has been a vexing question for more than ten years. Recently, based on early works [25], this question was addressed [26, 27] with the help of a novel 3-algebra. The theory based on seemingly the only unitary 3-algebra was found to be a theory for two M2-branes on an orbifold [28, 29]. More interestingly it could be rewritten as a super-Chern-Simons theory of two gauge groups with opposite level[30]. This fact, combined with the earlier attempts of constructing three-dimensional super-conformal field theory as Chern-Simons theory [31, 32], inspired people to search for the Chern-Simons-matter theory with higher supersymmetry. In [33], the one with $\mathcal{N} = 4$ supersymmetry

has been constructed. Very recently, a $\mathcal{N} = 6$ super Chern-Simons-matter theory was proposed in [34] to describe arbitrary number of M2-branes on the orbifold C^4/Z_k (For some other discussions on three-dimension $\mathcal{N} = 4, 5, 6$ super Chern-Simons-matter theories, see [35]-[39].) . In [34], the dual M/string description of large N limit of three-dimensional conformal theory was discussed. Depending on different values of the parameters in the field theory, this so-called ABJM theory can be dual to M-theory on $AdS_4 \times S^7/Z_k$ or IIA theory on $AdS_4 \times CP^3$.

The subject of this paper is to study the supersymmetric Wilson loops in the ABJM theory. The pure Chern-Simons theory is a topological theory and the correlation functions of Wilson loops in this theory give various knot invariants [40]. The ABJM theory includes matters and is no more a topological theory. We construct 1/6-BPS Wilson loops which involve the matter fields. Both the dimensional considerations and the supersymmetries lead us to include the product of two scalar fields in these operators.

As a first step to study these BPS Wilson loops, we perform the perturbative computations up to the order of λ^2 in the planar limit. On the other hand, we find a fundamental string solution ending on a circular loop on the boundary. The dual fundamental string solution here is quite similar to the one in AdS_5 case. This leads us to expect that this $F1$ -string solution is half-BPS, and it should be dual to a half-BPS Wilson loop which is to be constructed.

The other part of this paper is organized as following: in the next section, we construct some supersymmetric Wilson loops in ABJM theory and discuss the supersymmetries they preserve. In section 3, we first perform the perturbative calculations at the first two orders. In section 4, we will discuss the string theory dual of the to-be-constructed half-BPS Wilson loops. The conclusion and discussions are put in the last section. We also include some needed Feynman rules in Appendix A.

Note added

When this project was in progress, we learnt that this subject was also studied by Nadav Drukker, Jan Plefka and Donovan Young, whose paper [41] has overlap with ours.

2 Supersymmetric Wilson loops in super Chern-Simons-matter theory

In this paper, we study supersymmetric Wilson loop operators in three dimensional $\mathcal{N} = 6$ super Chern-Simons-matter theory proposed in [34]. The gauge group of this theory is $U_1(N) \times U_2(N)$. The matters include bifundamental superfields A_1, A_2 and anti-bifundamental superfields B_1, B_2 . The Chern-Simons part of the action is:

$$S_{CS} = \frac{k}{4\pi} \int \text{tr}(A_{(1)} \wedge dA_{(1)} + \frac{2}{3} A_{(1)}^3) - \text{tr}(A_{(2)} \wedge dA_{(2)} + \frac{2}{3} A_{(2)}^3). \quad (1)$$

Notice that the levels of these two group are opposite. The superpotential of this theory reads:

$$W = \frac{4\pi}{k} \text{Tr}(A_1 B_1 A_2 B_2 - A_1 B_2 A_2 B_1). \quad (2)$$

We denote $(A_1, A_2, \bar{B}_1, \bar{B}_2)$ as $Y^A, A = 1, \dots, 4$ which is in the fundamental representation of the R-symmetry group $SU_R(4)$.

The supersymmetry transformation of the gauge fields and the scalar fields are¹ [42]:

$$\delta A_\mu^{(1)} = -\frac{2\pi}{k} \left(Y^A \psi^{B\dagger} \gamma_\mu \omega_{AB} + \omega^{AB} \gamma_\mu \psi_A Y_B^\dagger \right), \quad (3)$$

$$\delta A_\mu^{(2)} = \frac{2\pi}{k} \left(\psi^{A\dagger} Y^B \gamma_\mu \omega_{AB} + \omega^{AB} \gamma_\mu Y_A^\dagger \psi_B \right), \quad (4)$$

$$\delta Y^A = i\omega^{AB} \psi_B, \quad (5)$$

$$\delta Y_A^\dagger = i\psi^{B\dagger} \omega_{AB}, \quad (6)$$

where ψ_A is the supersymmetry partner of Y^A and $\omega_{AB} = -\omega_{BA}$ are the supersymmetry parameters and they are 6 Majorana spinors in three dimensional spacetime. Here ω^{AB} is defined as

$$\omega^{AB} = \frac{1}{2} \epsilon^{ABCD} \omega_{CD}, \quad (7)$$

and we also have the following relation:

$$(\omega^{AB})_\alpha = ((\omega_{AB})_\alpha)^*. \quad (8)$$

¹We don't present the supersymmetry transformations of fermions, which are not essential to our study.

We will search for supersymmetric spacelike Wilson loop operators in the following class:

$$\begin{aligned} & \frac{1}{N} \text{Tr}_{U_1(N)} P \exp \left[i \int d\tau \left(A_\mu^{(1)} \dot{x}^\mu + i s_A^B(x) Y^A Y_B^\dagger \right) \right] \\ & \times \frac{1}{N} \text{Tr}_{U_2(N)} P \exp \left[i \int d\tau \left(A_\mu^{(2)} \dot{x}^\mu + i \tilde{s}_B^A(x) Y_A^\dagger Y^B \right) \right], \end{aligned} \quad (9)$$

where s_A^B and $\tilde{s}_B^A$ are Hermitian matrix, $A_\mu^{(1)} = A_{\mu a}^{(1)} (T^a)_i^j$, $A_\mu^{(2)} = A_{\mu \tilde{a}}^{(2)} (T^{\tilde{a}})_{\tilde{i}}^{\tilde{j}}$. We have also omitted the indices of Y^A , $Y_A^\dagger$ which are $(Y^A)_i^{\tilde{j}}$ and $(Y_A^\dagger)_{\tilde{i}}^j$. The indices i, j are the color indices of $U_1(N)$ and the indices $\tilde{i}, \tilde{j}$ are the color indices of $U_2(N)$. The normalization of the generators of the Lie algebra is chosen as $(T^a)_i^j (T^a)_k^l = \frac{1}{2} \delta_k^j \delta_i^l$, $(T^{\tilde{a}})_{\tilde{i}}^{\tilde{j}} (T^{\tilde{a}})_{\tilde{k}}^{\tilde{l}} = \frac{1}{2} \delta_{\tilde{k}}^{\tilde{j}} \delta_{\tilde{i}}^{\tilde{l}}$.

For the above Wilson line operators to be supersymmetric, the supersymmetry transformation of each factor should vanish by itself. The transformation of the first factor is proportional to

$$-\frac{2\pi}{k} (Y^A \psi^{B\dagger} \gamma_\mu \omega_{AB} + \omega^{AB} \gamma_\mu \psi_A Y_B^\dagger) \dot{x}^\mu - \omega^{AC} \psi_C Y_B^\dagger s_A^B - Y^A \psi^{C\dagger} \omega_{BC} s_A^B. \quad (10)$$

Then we get the following condition for the operator to be supersymmetric:

$$-\frac{2\pi}{k} \dot{x}^\mu \gamma_\mu \omega_{AB} - s_A^C \omega_{CB} = 0. \quad (11)$$

$$-\frac{2\pi}{k} \omega^{AB} \gamma_\mu \dot{x}^\mu - \omega^{CA} s_C^B = 0. \quad (12)$$

In fact, eq. (12) is complex conjugation of eq. (11). Notice that ω_{AB} is antisymmetric, we get from eq. (11):

$$-\frac{2\pi}{k} \dot{x}^\mu \gamma_\mu \omega_{BA} - s_A^C \omega_{BC} = 0, \quad (13)$$

from which and the following equation from relabelling the indices in eq. (11)

$$-\frac{2\pi}{k} \dot{x}^\mu \gamma_\mu \omega_{BA} - s_B^C \omega_{CA} = 0, \quad (14)$$

we have

$$s_A^C \omega_{BC} = s_B^C \omega_{CA}. \quad (15)$$

Since the general case is not easy to study, now we consider the special case in which s and $\tilde{s}$ are diagonal matrices:

$$s_A^B = s_A \delta_A^B, \quad \tilde{s}_A^B = \tilde{s}_A \delta_A^B. \quad (16)$$

In this case, from eq. (15), we get:

$$(s_A - s_B)\omega_{AB} = 0, \quad (17)$$

this is to say that if $s_A \neq s_B$, $\omega_{AB} = 0$.

Now we turn to consider the second factor:

$$\frac{1}{N} \text{Tr}_{U_2(N)} P \exp \left[i \int d\tau \left(A_\mu^{(2)} \dot{x}^\mu + i \tilde{s}_A Y_A^\dagger Y^A \right) \right]. \quad (18)$$

Similarly the supersymmetric condition for this factor gives:

$$\frac{2\pi}{k} \dot{x}^\mu \gamma_\mu \omega_{AB} - \tilde{s}_B \omega_{BA} = 0, \quad (19)$$

and its complex conjugation. As before, we can get

$$(\tilde{s}_A - \tilde{s}_B)\omega_{AB} = 0. \quad (20)$$

From eqs. (11) and (19-20), we get

$$(s_A - \tilde{s}_A)\omega_{AB} = 0. \quad (21)$$

So if $s_A \neq \tilde{s}_A$, ω_{AB} must be zero.

Using all of the above results, and also taking the following condition into account:

$$\omega_{AB}^* = \omega^{AB} = \frac{1}{2} \epsilon^{ABCD} \omega_{CD}, \quad (22)$$

and the γ -matrices in [43, 42], we find the 1/6-BPS Wilson loops which is defined by $s_{1,2} = \tilde{s}_{1,2} = -s_{3,4} = -\tilde{s}_{3,4} = \pm \frac{2\pi}{k} |\dot{x}|$. Choose the positive sign for s_1 , the Wilson loop is

$$\begin{aligned} & \frac{1}{N} \text{Tr}_{U_1(N)} P \exp \left(i \int d\tau (A_\mu^{(1)} \dot{x}^\mu + i \frac{2\pi}{k} (A_i \bar{A}_i - \bar{B}_i B_i) |\dot{\vec{x}}|) \right) \\ & \times \frac{1}{N} \text{Tr}_{U_2(N)} P \exp \left(i \int d\tau (A_\mu^{(2)} \dot{x}^\mu + i \frac{2\pi}{k} (\bar{A}_i A_i - B_i \bar{B}_i) |\dot{\vec{x}}|) \right). \end{aligned} \quad (23)$$

Here for straight Wilson line, the preserved supercharges are global.

If we put the spacelike Wilson line along the x^2 direction, the above discussions give that the unbroken supersymmetries $(Q^{AB})^\alpha$ are $(Q^{12})^2$ and $(Q^{34})^1$. They satisfy the following superalgebra:

$$\{(Q^{12})^2, (Q^{34})^1\} = -P_2, \{(Q^{12})^2, (Q^{12})^2\} = \{(Q^{34})^1, (Q^{34})^1\} = 0. \quad (24)$$

Notices that among three P_μ 's, only P_2 appears in the right-hand side of superalgebra. This is as expected since only translation along x^2 direction is unbroken.

The Wilson loops preserve two supercharges in $\mathcal{N} = 2$ and $\mathcal{N} = 3$ Chern-Simons-matter theory were studied in [32]. Our 1/6-BPS operators are the generalization to ABJM theory of these operators. It is interesting to see if one can find other BPS Wilson loops within and/or out of the class eq. (9) we begin with.

3 Perturbative computations

In this section, we perform the perturbative computations of the vacuum expectation values of the above 1/6-BPS Wilson loops in the large N limit. Then the expansion parameter is $\lambda = N/k$. We will do the computation up to the order of λ^2 . We list some needed Feynman rules in Appendix A. And we choose the Landau gauge and use the regularization by dimensional reduction as done in [44]. In this regularization scheme, we first perform all tensor contractions in strict three dimensions:

$$\epsilon_{\mu\nu\rho}\epsilon^{\mu\sigma\lambda} = \delta_\nu^\sigma\delta_\rho^\lambda - \delta_\nu^\lambda\delta_\rho^\sigma, \delta_\mu^\mu = 3, \quad (25)$$

then we regular the divergent scalar integrals by the dimensional continuation with $D = 3 - \epsilon$.

We will mainly focus on the circular Wilson loops. We parameterize the circle as

$$x^\mu(\tau) = (L \cos \tau, L \sin \tau, 0), \quad 0 \leq \tau \leq 2\pi, \quad (26)$$

with the x^3 direction being the time direction.

3.1 The calculations at the order of λ

We show in this subsection that the contribution at this order is zero. Due to the factor $\frac{2\pi}{k}$ in front of the product of two scalars, we need only consider the




Figure 1: The only contributing diagram at the order of λ .

contributions from the gauge field. There is only one contributing diagram as in Fig. 1. The contributions from this diagram reads:

$$\frac{i^2}{N} \int_0^{2\pi} d\tau_1 \int_0^{\tau_1} d\tau_2 \langle A^{\mu_1}(x(\tau_1)) A^{\mu_2}(x(\tau_2)) \rangle \frac{dx_{\mu_1}(\tau_1)}{d\tau_1} \frac{dx_{\mu_2}(\tau_2)}{d\tau_2}, \quad (27)$$

where the correlator of two gauge field is the free field one. The momentum space gluon propagator in the Landau gauge is proportional to ²

$$\frac{\epsilon^{\mu\nu\lambda} p_\lambda}{p^2}. \quad (28)$$

By Fourier transformation, we get the position space propagator

$$\langle A^{\mu_1}(x(\tau_1)) A^{\mu_2}(x(\tau_2)) \rangle \propto \frac{\epsilon^{\mu_1\mu_2\lambda} (x_1 - x_2)_\lambda}{|x_1 - x_2|^3}, \quad (29)$$

where $x_i = x(\tau_i)$, $i = 1, 2$. Using this result, and eq. (26), one can easily see that eq. (27) is equal to zero.

3.2 The calculations at the order of λ^2

We first focus on the contribution due to the expansion of the first factor. The planar diagrams at the order of λ^2 are listed in Fig. 2. From the discussions in the previous subsection, we know that the contribution from Figs. 2.a vanishes. The contribution from Fig. 2.f also vanishes due to the relative minus sign between $A_i \bar{A}_i$ and $\bar{B}_i B_i$ in the Wilson loop. The contribution

²We don't need to consider the color factor in the computations at this order.




Figure 2: The contributing diagrams at the order of λ^2 .

from Fig 2.b can be computed first in momentum space, then the result can be transferred to the position space by a Fourier transformation. The contribution from this diagram is proportional to

$$\frac{i^2}{Nk^2} \int_0^{2\pi} d\tau_1 \int_0^{\tau_1} d\tau_2 \dot{x}(\tau_1)_\mu |\dot{x}(\tau_2)| \int \frac{d^2 p}{(2\pi^3)} e^{[ip \cdot (x(\tau_1) - x(\tau_2))]} \frac{\epsilon^{\mu\nu\lambda} p_\nu}{p^2} \int \frac{d^3 q}{(2\pi)^3} \frac{2q_\lambda - p_\lambda}{q^2(p - q)^2}. \quad (30)$$

Since $\epsilon^{\mu\nu\lambda} p_\nu p_\lambda = 0$, we need only consider

$$\int \frac{d^3 q}{(2\pi)^3} \frac{q_\lambda}{q^2(p - q)^2} \quad (31)$$

in the last integration. From the Lorentz structure we know that this must be proportional to p_λ , so this will not contribute, either. We can check the above claim through direct computations using regularization by dimensional reduction in [44]:

$$\int \frac{d^D q}{(2\pi)^D} \frac{q_\lambda}{q^2(p - q)^2} \quad (32)$$

$$= \int \frac{d^D q}{(2\pi)^D} \int_0^1 dx \frac{q_\lambda}{[(1-x)q^2 + x(p-q)^2]^2} \quad (33)$$

$$= \int_0^1 dx \frac{d^D q}{(2\pi)^D} \frac{q_\lambda}{[(q-xp)^2 + (x-x^2)p^2]^2} \quad (34)$$

After the shift $q \rightarrow q + xp$, we get

$$\int_0^1 dx \frac{d^D q}{(2\pi)^D} \frac{q_\lambda + xp_\lambda}{[q^2 + (x-x^2)p^2]^2} \quad (35)$$

the first term vanish and the second term is proportional to p_λ , this confirms our claim.




Figure 3: One-matter-loop correction to the gluon propagator.

Now we turn to the contributions from Fig. 2.c-2.d. The contributions from Figs. 2.c are from the one-loop corrections to gluon propagator, and the contribution from Fig. 2.d includes two scalar propagators. The combination of these two parts has a quite remarkable structure, which is the same as the sum of the gluon propagator in the Landau gauge and the scalar propagator in the four dimensional $\mathcal{N} = 4$ super Yang-Mills theory.

The sum of the contributions from these diagrams is equal to:

$$\begin{aligned} & \frac{i^2}{N} \int_0^{2\pi} d\tau_1 \int_0^{\tau_1} d\tau_2 \left(\dot{x}(\tau_1)^\mu \dot{x}(\tau_2)^\nu \langle A^\mu(x(\tau_1))_i^j A^\nu(x(\tau_2))_j^k \rangle_{ol} - \left(\frac{2\pi}{k} \right)^2 \right. \\ & \times \left. |\dot{x}(\tau_1)| |\dot{x}(\tau_2)| \langle ((A_i \bar{A}_i - \bar{B}_i B_i)(x(\tau_1)))_i^j ((A_i \bar{A}_i - \bar{B}_i B_i)(x(\tau_2)))_j^k \rangle \right), \end{aligned} \quad (36)$$

where the subscript 'ol' means one-loop. The contributions from the gluon loop and the ghost loop cancel each other [44], noticing that we are using the regularization by dimensional reduction. The contributions from one-matter-loop are from Figs. 3.a-3.c. Both the one-fermion-loop and the one-scalar-loop correction to the gluon self-energy are [44] (see also [32])

$$\Pi_{\mu\nu}(p) = -\frac{N\delta_i^k}{64} \left(p\eta_{\mu\nu} - \frac{p_\mu p_\nu}{p} \right). \quad (37)$$

Taking the two gluon propagators into account, we get:

$$\begin{aligned} & \left(\frac{4\pi}{k}\right)^2 \frac{\epsilon^{\mu\mu_1\mu_2} p_{\mu_2}}{p^2} \Pi_{\mu_1\nu_1}(p) \frac{\epsilon^{\nu\nu_1\nu_2} p_{\nu_2}}{p^2} \\ &= -\frac{N\pi^2}{4k^2} \delta_i^k \frac{p^2 \eta^{\mu\nu} - p^\mu p^\nu}{p^3}. \end{aligned} \quad (38)$$

In the ABJM theory, we have $4N$ scalars and $4N$ fermions, so totally we get:

$$-\frac{2\pi^2 N^2}{k^2} \delta^{ab} \frac{p^2 \eta^{\mu\nu} - p^\mu p^\nu}{p^3}. \quad (39)$$

The second term of the above propagator will not contribute to the final answer due to the gauge invariance. In fact, the Fourier transformation of this term is proportional to

$$\partial_\mu^1 \partial_\nu^2 \log |x_1 - x_2|, \quad (40)$$

where $x_i = x(\tau_i)$, $\partial_\mu^i = \frac{\partial}{\partial x_i^\mu}$, $i = 1, 2$. So its contribution to eq. (36) is proportional to

$$\int_0^{2\pi} d\tau_1 \int_0^{\tau_1} d\tau_2 \dot{x}^\mu(\tau_1) \dot{x}^\nu(\tau_2) \partial_\mu^1 \partial_\nu^2 \log |x_1 - x_2|, \quad (41)$$

now we regularize the above integration as:

$$\int_0^{2\pi} d\tau_1 \dot{x}^\mu(\tau_1) \int_0^{\tau(\tilde{\epsilon})} d\tau_2 \frac{d}{d\tau_2} (\partial_\mu^1 \log |x_1 - x_2|), \quad (42)$$

where $\tau(\tilde{\epsilon}) = \tau_1 - \tilde{\epsilon}$ and $\tilde{\epsilon}$ is a regularization parameter. Then we get

$$\int_0^{2\pi} d\tau_1 \dot{x}^\mu(\tau_1) (\partial_\mu^1 \log |x_1 - x(\tau(\tilde{\epsilon}))| - \partial_\mu^1 \log |x_1 - x(0)|) \quad (43)$$

Using $\partial_\mu^1 \log |x_1 - y| \propto x_\mu(\tau_1)$ and $x(\tau_1) \cdot \dot{x}(\tau_1) = 0$, we get that eq. (2) is zero even before we take $\tilde{\epsilon} \rightarrow 0$. So this term doesn't contribute as claimed.

By performing a Fourier transformation of the first term in eq. (39) to position space, we get

$$\int \frac{d^3 p}{(2\pi)^3} e^{ip \cdot (x-y)} \left(-\frac{2\pi^2 N^2}{k^2} \delta_i^k \frac{1}{p^3} \right) = \frac{N^2 \eta^{\mu\nu} \delta^{ab}}{k^2 (x-y)^2}. \quad (44)$$

So the contribution to

$$i^2 \dot{x}_\mu \dot{y}_\nu \langle (A^\mu(x))_i^j (A^\nu(y))_j^k \rangle \quad (45)$$

at this order equals to

$$-\frac{N^2}{k^2} \frac{\dot{x} \cdot \dot{y}}{(x-y)^2} \delta_i^k, \quad (46)$$

where $x \equiv x_1, y \equiv x_2$. Now we turn to discuss the contribution due to the product of two scalar propagators which appears in Fig. 2.d. Using the position space propagator of (anti-)bifundamental scalars

$$\frac{1}{4\pi} \frac{1}{|x-y|} \delta_i^l \delta_k^{\tilde{j}}, \quad (47)$$

we get that

$$-\left(i\frac{2\pi}{k}\right)^2 |\dot{x}||\dot{y}| \langle (A_i \bar{A}_i - B_i \bar{B}_i)_i^j(x) (A_j \bar{A}_j - B_j \bar{B}_j)_j^k(y) \rangle = \frac{N^2}{k^2} \frac{|\dot{x}||\dot{y}|}{|x-y|^2} \delta_i^k. \quad (48)$$

The sum of eqs. (46) and (48) equals to

$$-\frac{N^2}{k^2} \frac{\dot{x} \cdot \dot{y} - |\dot{x}||\dot{y}|}{|x-y|^2} \delta_i^k. \quad (49)$$

When $x(\tau)$ is among the circle in eq. (26), the above result equals to

$$\frac{N^2}{2k^2} \delta_i^k \quad (50)$$

which is independent of the positions.

From this we get that eq. (36) is equal to

$$\frac{N^2}{2k^2} \int_0^{2\pi} d\tau_1 \int_0^{\tau_1} d\tau_2 = \frac{\pi^2 N^2}{k^2}. \quad (51)$$

The second factor of the Wilson loop gives the same contribution. Then the total contribution is $\frac{2\pi^2 N^2}{k^2}$.

Now at the order of λ^2 , we are left with the contribution from Fig. 2.e. This diagram comes from the pure Chern-Simons theory and was studied in details in [45]. The result from each factor in the Wilson line is $-\pi^2 N^2/(6k^2)$. So totally the contributions at the order of λ^2 is

$$2\pi^2 \lambda^2 - 2 \cdot \frac{\pi^2 \lambda^2}{6} = \frac{5}{3} \pi^2 \lambda^2. \quad (52)$$

4 Holographic description

Depending on the values of N and k , the gravity dual of the ABJM theory can be M-theory on $AdS_4 \times S^7/Z_k$ or type IIA theory on $AdS_4 \times CP^3$ with two-form and four-form fluxes. In this section we will focus on the type IIA case which corresponds to the situation when $k \ll N \ll k^5$. As the four dimensional super Yang-Mills case, the Wilson loop in the fundamental representation of the gauge group should be dual to a fundamental string in this background while the better description for the Wilson loop in higher dimensional representation should be higher dimensional probe D-branes or bubbling geometry.

In this section, we find a fundamental string solution ending on a circle on the boundary. The dual IIA string theory background is (We take $\alpha' = 1$):

$$ds_{string}^2 = \frac{R^3}{k} \left(\frac{1}{4} ds_{AdS_4}^2 + ds_{CP^3}^2 \right), \quad (53)$$

$$e^{2\phi} = \frac{R^3}{k^3}, \quad (54)$$

$$F_4 = \frac{3}{8} R^3 \epsilon_{AdS_4}, \quad (55)$$

$$F_2 = kJ, \quad (56)$$

where ϵ_{AdS_4} is the volume form of unit AdS_4 and J is the Kahler form of CP^3 , and R satisfies

$$R^3 = 2^{5/2} \pi \sqrt{Nk}. \quad (57)$$

Using the Poincare coordinates in the AdS_4 part, the metric can be written as

$$ds^2 = \frac{R^3}{4k} \frac{1}{y^2} (dy^2 + dx_1^2 + dx_2^2 + dx_3^2) + \frac{R^3}{k} ds_{CP^3}^2, \quad (58)$$

where $y = 0$ is the conformal boundary of AdS_4 . We put the circular Wilson loop at $x_1^2 + x_2^2 = R^2$ and $x_3 = 0$ on the boundary. As in the AdS_5 case [8, 9], the worldsheet of this fundamental string should be completely embedded in AdS_4 ending on the Wilson loop [34]. The Nambu-Goto action of the fundamental string is

$$S_{NG} = \frac{1}{2\pi} \int d\sigma d\tau \sqrt{\det \tilde{g}}, \quad (59)$$

where $\tilde{g}$ is the induced metric on the worldsheet. The fundamental string solution we are looking for is the same as the AdS_5 case [10, 11]:

$$x_1^2 + x_2^2 + y^2 = R^2. \quad (60)$$

The total action of this string is equal to

$$-\frac{R^3}{4k} = -\pi\sqrt{2\lambda}. \quad (61)$$

Since this fundamental string solution is quite similar to the one in IIB theory on $AdS_5 \times S^5$, we expect that this solution is half-BPS. It should be very interesting to construct the dual half-BPS Wilson loop in the field theory side.

5 Conclusion and Discussions

In this paper, we studied the supersymmetric Wilson loops in ABJM theory from both sides of the AdS_4/CFT_3 correspondence. In the field theory side, we constructed 1/6-BPS Wilson loop operators and perform the perturbative computations of their VEVs at the first two orders. In the string theory side, we found the fundamental string solution which could be dual to half-BPS circular Wilson loop.

There are a lot of interesting open questions we left for further studies. Here we list some of them. One natural question is to search for other BPS Wilson loops, especially the half-BPS one which seems to us not easy to construct. Even for the 1/6-BPS Wilson loops in this paper, our understanding is quite limited. Recently a class of Wilson loops operators are constructed in ABJM theory based on M-theory considerations [46] following the line in [9]. It is interesting to check if the 1/6-BPS Wilson loops belong to that class of operators and search for new supersymmetric Wilson loops in that class.

We also notice that Berkovits' pure string formalism [47, 48] has been used in the string theory dual of ABJM theory [49, 50, 51]. Based on these works, maybe one can study the Wilson loops in ABJM theory using the ideas in [52] for the AdS_5 case.

The fundamental string dual to half-BPS operator is almost the same as the one in AdS_5 case. However the string solution dual to the 1/6-BPS operators may also be embedded in the CP^3 part of the background in a non-trivial manner as the string solution in $AdS_5 \times S^5$ dual to 1/4-BPS

Wilson loop in super Yang-Mills theory [18]. It is interesting to explicitly find this solution. The probe brane dual to Wilson loops in higher dimensional representations need further studies as well. Another interesting question is to find $M2$ -brane in $AdS_4 \times S^7/Z_k$ dual to these Wilson loops.

Some membrane solutions in $AdS_4 \times S^7$ is found in [53, 54]. By generalizing them to membrane solutions $AdS_4 \times S^7/Z_k$ and using the reduction from M-theory to IIA theory, it is reasonable to get some fundamental string solution in $AdS_4 \times CP^3$. The field theory dual of these string solutions demands further studies.

Acknowledgments

The work was partially supported by NSFC Grant No.10535060,10775002 and NKBRPC (No. 2006CB805905). JB would like to thank Riccardo Argurio, Matteo Bertolini, Giulio Bonelli, Nadav Druker, Edi Gava, Chethan Gowdigere, Qing-Guo Huang, Kimyeong Lee, Juan Maldacena, Jan Plefka, Houman Safaai, Takao Suyama, Mahdi Torabian and Ho-Ung Yee for helpful discussions. We also thank Nadav Druker and Jan Plefka for correspondence prior to submission. The Feynman diagrams are drawn using Axodraw [55].

A The Feynman rules




Figure 4: Some needed Feynman rules

In this appendix, we list some of the Feynman rules needed in our perturbative calculation in ABJM theory. The diagrams are put in Fig. 6, and the corresponding expressions are the following:

1. Gluon propagator

$$\pm \delta^{ab} \frac{4\pi}{k} \epsilon^{\mu\nu\lambda} p_\lambda \frac{1}{p^2} \quad (62)$$

in the Landau gauge, where the up/down sign is corresponding to the first/sector factor of the gauge group.

2. Scalar propagator: $\frac{i}{p^2}$.
3. Fermion propagator: $\frac{ip_\mu \gamma^\mu}{p^2}$.
4. Three-gluon vertex: $\pm \frac{k}{4\pi} f^{abc} \epsilon^{\mu\nu\lambda}$. The up/down sign is corresponding to the first/sector factor of the gauge group.
5. Gluon-scalar-scalar vertex: $-i(p + p')^\mu (T^a)_{ij}$.
6. gluon-gluon-scalar-scalar vertex: $= i\eta^{\mu\nu} (\{T_a, T_b\})_{ij}$
7. Gluon-fermion-fermion vertex: $i\gamma^\mu (T^a)$.
8. One gluon emitted the Wilson loop: i .
9. Two scalars emitted the Wilson loop: $\pm i \frac{2\pi}{k}$ for A 's/ B 's.

The last two vertices are from the Wilson loop operator (we omit the color factors in the expressions of these two vertices). We don't give the explicit express from the six-scalar vertex and scalar-scalar-fermion-fermion vertex. We also omit the ghost propagator and gluon-ghost-ghost vertex.

References

- [1] J.M.Maldacena, “*The large N limit of superconformal field theories and supergravity*,” Adv. Theor. Math. Phys. **2**, 231 (1998) [hep-th/9711200].
- [2] S. S. Gubser, I. R. Klebanov and A. M. Polyakov, “*Gauge theory correlators from non-critical string theory*,” Phys. Lett. B **428**, 105 (1998) [arXiv:hep-th/9802109].
- [3] E. Witten, “*Anti-de Sitter space and holography*,” Adv. Theor. Math. Phys. **2**, 253 (1998) [arXiv:hep-th/9802150].

- [4] O. Aharony, S. S. Gubser, J. M. Maldacena, H. Ooguri and Y. Oz, “*Large N field theories, string theory and gravity*,” Phys. Rept. **323**, 183 (2000) [arXiv:hep-th/9905111].
- [5] J. K. Erickson, G. W. Semenoff and K. Zarembo, “*Wilson loops in $N = 4$ supersymmetric Yang-Mills theory*,” Nucl. Phys. **B582** (2000) 155–175 [hep-th/0003055].
- [6] N. Drukker and D. J. Gross, “*An exact prediction of $N = 4$ SUSYM theory for string theory*,” J. Math. Phys. **42** (2001) 2896–2914 [hep-th/0010274].
- [7] V. Pestun, “*Localization of gauge theory on a four-sphere and supersymmetric Wilson loops*,” [arXiv:0712.2824 [hep-th]].
- [8] S. J. Rey and J. T. Yee, “*Macroscopic strings as heavy quarks in large N gauge theory and anti-de Sitter supergravity*,” Eur. Phys. J. C **22**, 379 (2001) [arXiv:hep-th/9803001].
- [9] J. M. Maldacena, “*Wilson loops in large N field theories*,” Phys. Rev. Lett. **80**, 4859 (1998) [arXiv:hep-th/9803002].
- [10] D. E. Berenstein, R. Corrado, W. Fischler and J. M. Maldacena, “*The operator product expansion for Wilson loops and surfaces in the large N limit*,” Phys. Rev. D **59**, 105023 (1999) [arXiv:hep-th/9809188].
- [11] N. Drukker, D. J. Gross and H. Ooguri, “*Wilson loops and minimal surfaces*,” Phys. Rev. D **60**, 125006 (1999) [arXiv:hep-th/9904191].
- [12] N. Drukker and B. Fiol, “*All-genus calculation of Wilson loops using D -branes*,” JHEP **0502**, 010 (2005) [hep-th/0501109].
- [13] S. Yamaguchi, “*Wilson Loops of Anti-symmetric Representation and $D5$ -branes*,” JHEP **0605**, 037 (2006) [hep-th/0603208].
- [14] J. Gomis and F. Passerini, “*Holographic Wilson loops*,” JHEP **0608**, 074 (2006) [arXiv:hep-th/0604007].
- [15] J. Gomis and F. Passerini, “*Wilson loops as $D3$ -branes*,” JHEP **0701**, 097 (2007) [arXiv:hep-th/0612022].

- [16] B. Chen and W. He, “On $1/2$ -BPS Wilson-’t Hooft loops,” Phys.Rev.D74:126008,2006, [arXiv:hep-th/0607024].
- [17] K. Zarembo, “Supersymmetric Wilson loops,” Nucl. Phys. B **643**, 157 (2002) [arXiv:hep-th/0205160].
- [18] N. Drukker, “ $1/4$ BPS circular loops, unstable world-sheet instantons and the matrix model,” JHEP **0609**, 004 (2006) [arXiv:hep-th/0605151].
- [19] N. Drukker, S. Giombi, R. Ricci and D. Trancanelli, “On the $D3$ -brane description of some $1/4$ BPS Wilson loops,” JHEP **0704**, 008 (2007) [arXiv:hep-th/0612168].
- [20] N. Drukker, S. Giombi, R. Ricci and D. Trancanelli, “More supersymmetric Wilson loops,” Phys. Rev. D **76**, 107703 (2007), [arXiv:0704.2237 [hep-th]].
- [21] N. Drukker, S. Giombi, R. Ricci and D. Trancanelli, “Wilson loops: From four-dimensional SYM to two-dimensional YM,” Phys. Rev. D **77**, 047901 (2008) [arXiv:0707.2699 [hep-th]].
- [22] N. Drukker, S. Giombi, R. Ricci and D. Trancanelli, “Supersymmetric Wilson loops on S^3 ,” JHEP **0805**, 017 (2008) [arXiv:0711.3226 [hep-th]].
- [23] A. Bassetto, L. Griguolo, F. Pucci and D. Seminara, “Supersymmetric Wilson loops at two loops,” JHEP **0806**, 083 (2008) [arXiv:0804.3973 [hep-th]].
- [24] D. Young, “BPS Wilson Loops on S^2 at Higher Loops,” JHEP **0805**, 077 (2008) [arXiv:0804.4098 [hep-th]].
- [25] A. Basu and J. A. Harvey, “The $M2$ - $M5$ brane system and a generalized Nahm’s equation,” Nucl. Phys. B **713**, 136 (2005) [arXiv:hep-th/0412310].
- [26] J. Bagger and N. Lambert, “Modeling multiple $M2$ ’s”, Phys. Rev. D**75**,045020(2007)[arXiv:hep-th/0611108].
“Gauge Symmetry and Supersymmetry of Multiple $M2$ -Branes”, Phys. Rev. D**77**, 065008(2008) [arXiv:0711.0955]. “Comments on Multiple $M2$ -branes”, JHEP **0802**, 105(2008)[arXiv:0712.3738].

- [27] A. Gustavsson, “*Algebraic structures on parallel M2-branes*”, arXiv:0709.1260.
“*One-loop corrections to Bagger-Lambert theory*”, arXiv:0805.4443.
- [28] J. Distler et.al., “*M2-branes on M-folds*”, JHEP **0805**, 038(2008)[arXiv:0804.1256].
- [29] N. Lambert and D. Tong, “*Membranes on an Orbifold*,” Phys. Rev. Lett. **101**, 041602 (2008) [arXiv:0804.1114 [hep-th]].
- [30] M. Van Raamsdonk, “*Comments on the Bagger-Lambert theory and multiple M2-branes*”, arXiv: 0803.3803[hep-th].
M.A. Bandres, A.E. Lipstein and J.H. Schwarz, “ *$N = 8$ Superconformal Chern-Simons Theories*,”, arXiv:0803.3242[hep-th].
- [31] J. H. Schwarz, “*Superconformal Chern-Simons theories*,” JHEP **0411**, 078 (2004) [arXiv:hep-th/0411077].
- [32] D. Gaiotto and X. Yin, “*Notes on superconformal Chern-Simons-matter theories*,” JHEP **0708**, 056 (2007) [arXiv:0704.3740 [hep-th]].
- [33] D. Gaiotto and E. Witten, “*Janus Configurations, Chern-Simons Couplings, and the θ -angle in $\mathcal{N} = 4$ super-Yang-Mills theory*”, arXiv:0804.2907[hep-th].
- [34] O. Aharony, O. Bergman, D. L. Jafferis and J. Maldacena, “ *$N=6$ superconformal Chern-Simons-matter theories, M2-branes and their gravity duals*,” arXiv:0806.1218 [hep-th].
- [35] K. Hosomichi, K. M. Lee, S. Lee, S. Lee and J. Park, “ *$N=4$ Superconformal Chern-Simons Theories with Hyper and Twisted Hyper Multiplets*,” JHEP **0807**, 091 (2008) [arXiv:0805.3662 [hep-th]].
- [36] K. Hosomichi, K. M. Lee, S. Lee, S. Lee and J. Park, “ *$N=5,6$ Superconformal Chern-Simons Theories and M2-branes on Orbifolds*,” arXiv:0806.4977 [hep-th].
- [37] M. Schnabl and Y. Tachikawa, “*Classification of $N=6$ superconformal theories of ABJM type*,” arXiv:0807.1102 [hep-th].

- [38] E. A. Bergshoeff, O. Hohm, D. Roest, H. Samtleben and E. Sezgin, “*The Superconformal Gaugings in Three Dimensions*,” arXiv:0807.2841 [hep-th].
- [39] O. Aharony, O. Bergman and D. L. Jafferis, “*Fractional M2-branes*,” arXiv: 0807.4924 [hep-th].
- [40] E. Witten, “*Quantum field theory and the Jones polynomial*,” Commun. Math. Phys. **121**, 351 (1989).
- [41] N. Drukker, J. Plefka and D. Young, “*Wilson loops in 3-dimensional $\mathcal{N} = 6$ supersymmetric Chern-Simons and their string theory dual*,” arXiv: 0809.2787 [hep-th].
- [42] S. Terashima, “*On M5-branes in N=6 Membrane Action*,” arXiv:0807.0197 [hep-th].
- [43] M. Benna, I. Klebanov, T. Klose and M. Smedback, “*Superconformal Chern-Simons Theories and AdS_4/CFT_3 Correspondence*,” JHEP **0809**, 072 (2008) [arXiv:0806.1519 [hep-th]].
- [44] W. Chen, G. W. Semenoff and Y. S. Wu, “*Two loop analysis of nonAbelian Chern-Simons theory*,” Phys. Rev. D **46**, 5521 (1992) [arXiv:hep-th/9209005].
- [45] E. Guadagnini, M. Martellini and M. Mintchev, “*Wilson Lines in Chern-Simons Theory and Link Invariants*,” Nucl. Phys. B **330**, 575 (1990).
- [46] D. Berenstein and D. Trancanelli, “*Three-dimensional N=6 SCFT’s and their membrane dynamics*,” arXiv:0808.2503 [hep-th].
- [47] N. Berkovits, “*Super-Poincare covariant quantization of the superstring*,” JHEP **0004**, 018 (2000) [arXiv:hep-th/0001035].
- [48] N. Berkovits, “*ICTP lectures on covariant quantization of the superstring*,” arXiv:hep-th/0209059.
- [49] P. Fre’ and P. A. Grassi, “*Pure Spinor Formalism for $Osp(N|4)$ backgrounds*,” arXiv:0807.0044 [hep-th].
- [50] G. Bonelli, P. A. Grassi and H. Safaai, “*Exploring Pure Spinor String Theory on $AdS_4 \times CP^3$* ,” arXiv:0808.1051 [hep-th].

- [51] R. D'Auria, P. Fre', P. A. Grassi and M. Trigiante, "*Superstrings on $AdS_4 \times CP^3$ from Supergravity*," arXiv:0808.1282 [hep-th].
- [52] G. Bonelli and H. Safaai, "*On gauge/string correspondence and mirror symmetry*," JHEP **0806**, 050 (2008) [arXiv:0804.2629 [hep-th]].
- [53] O. Lunin, "*1/2-BPS states in M theory and defects in the dual CFTs*," JHEP **0710**, 014 (2007) [arXiv:0704.3442 [hep-th]].
- [54] B. Chen, "*The Self-dual String Soliton in $AdS_4 \times S^7$ spacetime*," Eur. Phys. J. C **54**, 489 (2008) [arXiv:0710.2593 [hep-th]].
- [55] J. A. M. Vermaseren, "*Axodraw*," Comput. Phys. Commun. **83**, 45(1994).